

## WEEKLY NEW BRUNSWICK INFLUENZA REPORT

Reporting period: December 8 to December 14 2019 (week 50)

### Summary

#### In New Brunswick, influenza activity was low in week 50

##### New Brunswick:

- There have been 12 positive influenza cases in week 50. Since the beginning of the new season, 29 cases has been reported, 11 influenza A (H1N1)pdm09, 7 influenza A (H3), 1 influenza A (unsubtyped) and 10 influenza B.
- There has been 1 new influenza associated hospitalization during week 50. So far this season, 7 influenza associated hospitalizations have been reported and no deaths.
- The ILI consultation rate was 4.9 consultations per 1,000 patients visits in week 50. The ILI rate was below the expected levels for this time of year.
- One new ILI outbreak was reported in a school in week 50. So far this season, 1 influenza outbreak has been reported in a nursing home and 2 ILI outbreaks were reported in schools.

##### Canada:

- Influenza activity continued to increase across multiple indicators, with regions throughout the country reporting increased influenza activity.
- In week 50, the proportion of influenza B detections has increased resulting in an equal proportion of influenza A and B detections nationally. This level of influenza B activity is higher than average for this time of year and usually not seen until February or March.
- This season, the majority of hospitalizations reported by participating provinces and territories have been associated with influenza A(H3N2). Among sentinel pediatric hospitalizations with influenza, approximately 55% of cases were associated with influenza B.

##### International:

###### Seasonal influenza:

In the temperate zone of the northern hemisphere, respiratory illness indicators and influenza activity continued to increase in most countries. In the Caribbean and Central American countries, influenza activity was low overall, except for Cuba where increased detections of influenza B/Victoria lineage viruses were reported. In tropical South American countries, influenza activity remained low. In tropical Africa, influenza activity remained elevated in some countries of Middle and Western Africa. In Southern Asia, influenza activity was low across reporting countries, but was reported at high levels in the Islamic Republic of Iran in recent weeks. In South East Asia, influenza activity continued to be reported in Lao PDR and the Philippines. In the temperate zones of the southern hemisphere, influenza activity remained at inter-seasonal levels. Worldwide, seasonal influenza A(H3N2) viruses accounted for the majority of detections.

###### Emerging Respiratory Viruses:


- MERS CoV:
  - WHO: [http://www.who.int/csr/disease/coronavirus\\_infections/en/](http://www.who.int/csr/disease/coronavirus_infections/en/)
  - CDC: <http://www.cdc.gov/coronavirus/mers/>
  - Updated Risk Assessment (August 2018): [http://www.who.int/csr/disease/coronavirus\\_infections/risk-assessment-august-2018.pdf?ua=1](http://www.who.int/csr/disease/coronavirus_infections/risk-assessment-august-2018.pdf?ua=1)
- Avian Influenza:
  - WHO: [www.who.int/csr/disease/avian\\_influenza/en/index.html](http://www.who.int/csr/disease/avian_influenza/en/index.html)

### 1) Influenza Laboratory Data<sup>1</sup>

- Influenza activity was low in week 50.
- Twelve influenza cases were reported during week 50, 4 influenza A (H1N1)pdm09, 2 influenza A (H3), 1 influenza A (unsubtyped) and 5 influenza B.
- Since the beginning of the season, 29 influenza cases have been reported, 11 influenza A (H1N1)pdm09, 7 influenza A (H3), 1 influenza A (unsubtyped) and 10 influenza B.


<sup>1</sup> Surveillance specimens are submitted by recruited New Brunswick Sentinel Practitioner Influenza Network (NB SPIN) practitioners, which are comprised of sites in Emergency Rooms, in Family Practice, in First Nations communities, in Nursing Home, in Universities and in Community Health Centers. Diagnostic specimens are submitted by physicians in the community/hospital setting. Influenza laboratory data is comprised of results from surveillance and diagnostic specimens. All laboratory specimens are tested using a real-time PCR assay, which is a rapid detection method designed for detection of all known variants of influenza A and B. All laboratory-confirmed cases are reported for the week when laboratory confirmation was received.

Graph 1: Number and percent of positive influenza specimens<sup>2</sup> in New Brunswick by week, up to December 14, 2019 (data source: G. Dumont Lab results)



<sup>2</sup> Total number of positive influenza tests is higher than number of cases since some individuals had co-infection of A & B simultaneously.

Figure 2: Influenza/ILI activity levels<sup>3</sup> by Health Zones, in New Brunswick, for week 50, season 2019/2020.



<sup>3</sup> No activity is defined as no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported. Sporadic activity is defined as sporadically occurring ILI and lab confirmed influenza detection(s) with no outbreaks detected within the influenza surveillance region. Localized activity is defined as evidence of increased ILI with lab confirmed influenza detection(s) and outbreaks in schools, hospitals, residential institutions and/or other types of facilities occurring in less than 50% of the influenza surveillance region. Widespread activity is defined as evidence of increased ILI with lab confirmed influenza detection(s) and outbreaks in schools, hospitals, residential institutions and/or other types of facilities occurring in greater than or equal to 50% of the influenza surveillance region.


**Table 1:** Positive influenza cases by Health Region, in New Brunswick for reporting week, cumulative current and previous seasons.  
 (data source: G. Dumont lab results up to December 14, 2019)

Zone	Reporting period: December/08/2019–December/14/2019						Cumulative: (2019/2020 season) Aug./25/2019 –December/14/2019						Cumulative: (2018/2019 season) Aug./26/2018 –Aug./24/2019									
	A					B	A & B co- infection					A					B	A & B co- infectio n				
	A(H3)	(H1N1) pdm09	Unsubty ped/ Other	A Total	Total	Total	A(H3)	(H1N1) pdm09	Unsubty ped/ Other	A Total	Total	Total	(H3)	(H1N1) pdm09	Unsubty ped/ Other	A Total	Total	Total				
Zone 1	2	1	1	4	3	0	5	4	1	10	5	0	29	97	1163	1289	130	3				
Zone 2	0	1	0	1	1	0	1	1	0	2	2	0	6	47	293	346	58	0				
Zone 3	0	0	0	0	0	0	1	2	0	3	2	0	9	39	260	308	3	0				
Zone 4	0	0	0	0	0	0	0	0	0	0	0	0	2	28	135	165	6	0				
Zone 5	0	0	0	0	0	0	0	1	0	1	0	0	2	20	84	106	127	1				
Zone 6	0	1	0	1	0	0	0	2	0	2	0	0	5	36	200	241	14	0				
Zone 7	0	1	0	1	1	0	0	1	0	1	1	0	9	23	160	192	19	0				
<b>Total NB</b>	<b>2</b>	<b>4</b>	<b>1</b>	<b>7</b>	<b>5</b>	<b>0</b>	<b>7</b>	<b>11</b>	<b>1</b>	<b>19</b>	<b>10</b>	<b>0</b>	<b>62</b>	<b>290</b>	<b>2295</b>	<b>2647</b>	<b>357</b>	<b>4</b>				

2) ILI Consultation Rates<sup>4</sup>

- For week 50, the ILI consultation rate was 4.9 consultations per 1,000 patients visits. The ILI rate was below the expected levels for this time of year.
- During week 50, the sentinel response rate was 32%, for both the FluWatch sentinel physicians and the NB SPIN practitioners.

Graph 2: ILI Consultation Rates in New Brunswick, by report week, season 2019/20 compared to previous seasons\*



\* The mean rate was based on data from the 1996/97 to 2018/2019 seasons and excludes the Pandemic season (2009/10).


<sup>4</sup> A total of 28 practitioner sites (16 FluWatch sentinel physicians and 12 NB SPIN sites) are recruited this season to report the number of ILI patients and total patient consultations one day during a reporting week.

### 3) ILI and Laboratory-Confirmed Outbreak Data

Table 2: ILI activity/outbreaks in New Brunswick nursing homes and schools for the reporting week, current and previous seasons.

	Reporting period: December/08/2019-December/14/2019			Cumulative # of outbreaks season 2019-2020	Cumulative # of outbreaks season 2018-2019
	Lab-confirmed outbreaks in Nursing homes <sup>5</sup>	ILI school outbreaks <sup>6</sup>	Lab-confirmed outbreaks in Other settings <sup>4</sup>		
Zone 1	0 out of 13	0 out of 74	0	0	12
Zone 2	0 out of 16	0 out of 81	0	0	13
Zone 3	0 out of 14	0 out of 95	0	1	6
Zone 4	0 out of 6	0 out of 22	0	0	0
Zone 5	0 out of 2	0 out of 18	0	0	0
Zone 6	0 out of 9	0 out of 35	0	0	4
Zone 7	0 out of 4	1 out of 27	0	2	8
<b>Total NB</b>	<b>0 out of 64</b>	<b>1 out of 352</b>	<b>0</b>	<b>3</b>	<b>43</b>

Graph 3: Number of Influenza Outbreaks (nursing homes, hospitals, other) and ILI Outbreaks (schools) reported to Public Health in New Brunswick, by report week, season 2019/20.




<sup>5</sup> Two or more ILI cases within a seven day period, including at least one laboratory-confirmed case of influenza. Outbreaks are reported in the week when laboratory confirmation is received.

<sup>6</sup> Schools reporting greater than 10% absenteeism which is likely due to ILI.

#### 4) Influenza associated Hospitalization<sup>7</sup> and Death<sup>8</sup> Surveillance<sup>9</sup>

Graph 4: Influenza associated Hospitalizations and ICU admissions in New Brunswick, by week of hospitalization for current and past season.\*



\*Those who had been hospitalized 15 days or more prior to laboratory confirmation date were excluded from the graph

\*\*No deaths have been reported so far in season 2019-2020.

National Flu Watch Program - Additional information on influenza activity in Canada and around the world is available on the Public Health Agency of Canada's website at: <http://www.phac-aspc.gc.ca/fluwatch/>

#### Other Links:

World: [http://www.who.int/influenza/surveillance\\_monitoring/updates/latest\\_update\\_GIP\\_surveillance/en/index.html](http://www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/index.html)

Europe: [http://www.ecdc.europa.eu/en/healthtopics/seasonal\\_influenza/epidemiological\\_data/Pages/Weekly\\_Influenza\\_Surveillance\\_Overview.aspx](http://www.ecdc.europa.eu/en/healthtopics/seasonal_influenza/epidemiological_data/Pages/Weekly_Influenza_Surveillance_Overview.aspx)

PAHO: [http://new.paho.org/hq/index.php?option=com\\_content&task=blogcategory&id=805&Itemid=569](http://new.paho.org/hq/index.php?option=com_content&task=blogcategory&id=805&Itemid=569)

Australia: <http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-surveil-ozflu-flucurr.htm>

New Zealand: [http://www.surv.esr.cri.nz/virology/influenza\\_weekly\\_update.php](http://www.surv.esr.cri.nz/virology/influenza_weekly_update.php)

Argentina: <http://www.msal.gov.ar/>

South Africa: <http://www.nicd.ac.za/>

US: [www.cdc.gov/flu/weekly/](http://www.cdc.gov/flu/weekly/)

Prepared by the Communicable Disease Control Unit, Office of the Chief Medical Officer of Health, Tel: (506) 444-3044

<sup>7</sup> Hospitalizations (including ICU admissions) are influenza associated; they may or may not be due to influenza.

<sup>8</sup> Deaths are influenza associated; influenza may not be the direct cause of death.

<sup>9</sup> In early January 2014, the Office of the Chief Medical Officer of Health implemented a new provincial surveillance system in collaboration with the Regional Health Authorities to monitor influenza-associated hospitalizations, intensive care unit admissions and deaths. A standardized Enhanced Surveillance Form is used to collect data on hospitalizations.