

WEEKLY NEW BRUNSWICK INFLUENZA REPORT

Reporting period: February 21 to February 27, 2021 (week 8)

Summary

In New Brunswick, influenza activity remained at inter-seasonal levels in week 8

New Brunswick:

- There have been no positive influenza cases in week 8. Since the beginning of the season, 1 case of influenza B has been reported.
- There has been no new influenza associated hospitalizations during week 8. Since the beginning of the season, no hospitalizations have been reported and no deaths.
- The ILI consultation rate was 0.0 per 1,000 patients visits for week 8. The ILI rate was lower than the expected levels for this time of year.
- No influenza outbreaks were reported in week 8. So far this season, no influenza outbreaks have been reported.

Canada:

- All indicators of influenza activity remain exceptionally low for this time of year, despite continued monitoring for influenza across Canada.
- To date, there is no evidence of community circulation of influenza. Influenza testing continues at seasonal levels. No laboratory detections of influenza were reported in week 8. Influenza activity remains below the threshold required to declare the start of the 2020-21 influenza season.
- Two influenza-like-illness (ILI) outbreaks were reported in schools and/or daycares in week 8. To date this season, 117 ILI outbreaks have been reported in schools and/or daycares. No laboratory-confirmed outbreaks of influenza have been reported to date this season.
- 12,030 participants reported to FluWatchers and 12 (0.10%) participants reported cough and fever, which remains low compared to previous seasons.
- Influenza surveillance indicators may be influenced by the COVID-19 pandemic, including changes in healthcare-seeking behaviour, impacts of public health measures and influenza testing capacity. Current data should be interpreted with consideration to this context.

International:

Seasonal influenza:

The current influenza surveillance data should be interpreted with caution as the ongoing COVID-19 pandemic have influenced to varying extents health seeking behaviours, staffing/routines in sentinel sites, as well as testing priorities and capacities in Member States. The various hygiene and physical distancing measures implemented by Member States to reduce SARS-CoV-2 virus transmission have likely played a role in reducing influenza virus transmission. Globally, despite continued or even increased testing for influenza in some countries, influenza activity remained at lower levels than expected for this time of the year. In the temperate zone of the northern hemisphere, influenza activity remained below baseline, though sporadic detections of influenza A and B viruses were reported in some countries. In the temperate zones of the southern hemisphere, influenza activity was reported at inter-seasonal level. In the Caribbean and Central American countries, sporadic influenza detections were reported. Severe acute respiratory infection (SARI) activity was low in most reporting countries. In tropical South America, sporadic detections were reported in Colombia. In tropical Africa, influenza activity was reported in some reporting countries in Western, Middle and Eastern Africa in recent weeks. In Southern Asia, sporadic influenza detections were reported in India. In South East Asia, influenza A(H3N2) detections continued to be reported in most reporting countries. Worldwide, influenza B detections accounted for the majority of the very low numbers of detections reported.

Emerging Respiratory Viruses:


- **COVID-19:** On December 31, 2019, a cluster of cases of pneumonia was reported in Wuhan, China, and the cause has been confirmed as a new coronavirus that has not previously been identified in humans (COVID-19). As of March 9, 2021, 893,518 cases of COVID-19 infection in Canada have been identified with 22,304 deaths. One thousand four hundred and sixty cases have been identified in New Brunswick with 30 deaths. As of March 10, the WHO reported globally 117 332 262 confirmed cases and 2 605 356 deaths in approximately 223 countries/territories/areas. For more timely updates, please visit the following websites:
 - WHO: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>
 - PHAC: <https://www.canada.ca/en/public-health/services/diseases/2019-novel-coronavirus-infection.html>
 - NB : https://www2.gnb.ca/content/gnb/en/departments/ocmoh/cdc/content/respiratory_diseases/coronavirus.html
- **MERS CoV:**
 - WHO: http://www.who.int/csr/disease/coronavirus_infections/en/
 - CDC: <http://www.cdc.gov/coronavirus/mers/>

1) Influenza Laboratory Data¹

¹ Surveillance specimens are submitted by recruited New Brunswick Sentinel Practitioner Influenza Network (NB SPIN) practitioners, which are comprised of sites in Emergency Rooms, in Family Practice, in First Nations communities, in Nursing Home, in Universities and in Community Health Centers. Diagnostic specimens are submitted by physicians in the community/hospital setting. Influenza laboratory data is comprised of results from surveillance and diagnostic specimens. All laboratory specimens are tested using a real-time PCR assay, which is a rapid detection method designed for detection of all known variants of influenza A and B. All laboratory-confirmed cases are reported for the week when laboratory confirmation was received.

- Influenza activity remained at inter-seasonal levels in week 8.
- No influenza cases were reported during week 8.
- Since the beginning of the season, 1 influenza case has been reported, an influenza B virus².

Graph 1: Number and percent of positive influenza specimens³ in New Brunswick by week, up to February 27, 2021 (data source: G. Dumont Lab results)




*The increase in influenza laboratory tests seen between week 50 and week 5 is due to a change in testing process (co-testing for influenza and COVID)

² This positive influenza detection is associated with recent live attenuated influenza vaccine receipt and does not represent community circulation of seasonal influenza viruses.

³ Total number of positive influenza tests is higher than number of cases since some individuals had co-infection of A & B simultaneously.

Figure 2: Influenza/ILI activity levels⁴ by Health Zones, in New Brunswick, for week 8, season 2020/2021.



⁴ No activity is defined as no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported.

Sporadic activity is defined as sporadically occurring ILI and lab confirmed influenza detection(s) with no outbreaks detected within the influenza surveillance region.

Localized activity is defined as evidence of increased ILI with lab confirmed influenza detection(s) and outbreaks in schools, hospitals, residential institutions and/or other types of facilities occurring in less than 50% of the influenza surveillance region.

Widespread activity is defined as evidence of increased ILI with lab confirmed influenza detection(s) and outbreaks in schools, hospitals, residential institutions and/or other types of facilities occurring in greater than or equal to 50% of the influenza surveillance region.

Table 1: Positive influenza cases by Health Region, in New Brunswick for reporting week, cumulative current and previous seasons.
(data source: G. Dumont lab results up to February 27, 2021)


| Zone | Reporting period: February/21/2021–February/27/2021 | | | | | | Cumulative: (2020/2021 season) Aug./23/2020 –February/27/2021 | | | | | | Cumulative: (2019/2020 season) Aug./25/2019 –Aug./22/2020 | | | | | |
|-----------------|--|-----------------|--------------------------|------------|----------|---------------------------|--|-----------------|--------------------------|------------|-----------|---------------------------|---|-----------------|--------------------------|------------|-------------|-------------------------------|
| | A | | | | B | A & B co- infection | A | | | | B | A & B co- infection | A | | | | B | A & B co- infectio n |
| | A(H3) | (H1N1) pdm09 | Unsubty ped/ Other | A Total | Total | Total | A(H3) | (H1N1) pdm09 | Unsubty ped/ Other | A Total | Total | Total | (H3) | (H1N1) pdm09 | Unsubty ped/ Other | A Total | Total | Total |
| Zone 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1* | 0 | 9 | 28 | 324 | 361 | 665 | 3 |
| Zone 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 11 | 121 | 135 | 96 | 2 |
| Zone 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 8 | 102 | 111 | 188 | 5 |
| Zone 4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 7 | 43 | 51 | 212 | 1 |
| Zone 5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 10 | 5 | 85 | 100 | 17 | 1 |
| Zone 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 6 | 7 | 120 | 133 | 98 | 1 |
| Zone 7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 65 | 68 | 103 | 0 |
| Total NB | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1* | 0 | 30 | 69 | 860 | 959 | 1379 | 13 |

*This positive influenza detection is associated with recent live attenuated influenza vaccine receipt and does not represent community circulation of seasonal influenza viruses.

2) ILI Consultation Rates⁵

- For week 8, the ILI consultation rate was 0.0 consultations per 1,000 patients visits. The ILI rate was lower than the expected levels for this time of year.
- During week 8, the sentinel response rate was 21% for both the FluWatch sentinel physicians and the NB SPIN practitioners.

Graph 2: ILI Consultation Rates in New Brunswick, by report week, season 2020/21 compared to previous seasons*




3) ILI and Laboratory-Confirmed Outbreak Data

Table 2: New ILI activity/outbreaks in New Brunswick nursing homes and schools* for the reporting week and current season.

| | Reporting period: February/21/2021 to February/27/2021 | | | Cumulative # of outbreaks season 2020-2021* |
|-----------------|---|-------------------------------------|--|---|
| | Lab-confirmed outbreaks in Nursing homes ⁶ | ILI school outbreaks ⁷ * | Lab-confirmed outbreaks in Other settings ⁴ | |
| Zone 1 | 0 out of 15 | 0 out of 74 | 0 | 28 |
| Zone 2 | 0 out of 16 | 0 out of 81 | 0 | 14 |
| Zone 3 | 0 out of 16 | 0 out of 95 | 0 | 23 |
| Zone 4 | 0 out of 5 | 0 out of 22 | 0 | 0 |
| Zone 5 | 0 out of 2 | 0 out of 18 | 0 | 0 |
| Zone 6 | 0 out of 9 | 0 out of 35 | 0 | 0 |
| Zone 7 | 0 out of 5 | 0 out of 27 | 0 | 2 |
| Total NB | 0 out of 68 | 0 out of 352 | 0 | 67* |

*During this influenza season, 2020-2021, the number of ILI outbreaks in school (based on greater than 10% absenteeism in school due to ILI symptoms, which for many schools cannot be determined) will likely be skewed due to the COVID-19 pandemic, the prudence of parents/guardians to send their children to school and their interpretation of the home isolation requirements. Therefore, the number of ILI outbreaks in schools should be interpreted with caution and should not be compared to previous seasons.

Graph 3: Number of Influenza Outbreaks (nursing homes, hospitals, other) and ILI Outbreaks (schools) reported to Public Health in New Brunswick, by report week, season 2020/21.




⁶ Two or more ILI cases within a seven-day period, including at least one laboratory-confirmed case of influenza. Outbreaks are reported in the week when laboratory confirmation is received.

⁷ Schools reporting greater than 10% absenteeism which is likely due to ILI.

4) Influenza associated Hospitalization⁸ and Death⁹ Surveillance¹⁰

Graph 4: Influenza associated Hospitalizations and ICU admissions in New Brunswick, by week of hospitalization for current and past season.*



*Those who had been hospitalized 15 days or more prior to laboratory confirmation date were excluded from the graph

**No deaths have been reported so far in season 2020-2021.

National Flu Watch Program - Additional information on influenza activity in Canada and around the world is available on the Public Health Agency of Canada's website at: <http://www.phac-aspc.gc.ca/fluwatch/>

Other Links:

World: http://www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/index.html

Europe: http://www.ecdc.europa.eu/en/healthtopics/seasonal_influenza/epidemiological_data/Pages/Weekly_Influenza_Surveillance_Overview.aspx

PAHO: http://new.paho.org/hq/index.php?option=com_content&task=blogcategory&id=805&Itemid=569

Australia: <http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-surveil-ozflu-flucurr.htm>

New Zealand: http://www.surv.esr.cri.nz/virology/influenza_weekly_update.php

Argentina: <http://www.msal.gov.ar/>

South Africa: <http://www.nicd.ac.za/>

US: www.cdc.gov/flu/weekly/

Prepared by the Communicable Disease Control Unit, Office of the Chief Medical Officer of Health, Tel: (506) 444-3044

⁸ Hospitalizations (including ICU admissions) are influenza associated; they may or may not be due to influenza.

⁹ Deaths are influenza associated; influenza may not be the direct cause of death.

¹⁰ In early January 2014, the Office of the Chief Medical Officer of Health implemented a new provincial surveillance system in collaboration with the Regional Health Authorities to monitor influenza-associated hospitalizations, intensive care unit admissions and deaths. A standardized Enhanced Surveillance Form is used to collect data on hospitalizations.